



International
Animal Health Products
THE AUSTRALIAN COMPANY

M^{EC}WORMA[®] & BOT

BROAD-SPECTRUM WORM PASTE FOR HORSES OF ALL AGES

PRESENTATION: Oral paste in dial-a-dose syringe.

ACTIVE CONSTITUENTS: Each g contains:
ABAMECTIN 3.64 mg.

PROPERTIES:

FARNAM[®] M^{EC}WORMA[®] & BOT is highly effective (up to 100% activity) in the control and treatment of large and small strongyles (*Strongylus* and *Cyathostomum* spp.) including arterial larval stages and benzimidazole resistant strains, bots (*Gasterophilus* spp.), cutaneous onchocerciasis and nematodes including *Habronema* and *Draschia* spp. which cause skin lesions (summer sores).

M^{EC}WORMA & BOT contains abamectin, which is the only natural form of the macrocyclic lactone group of anthelmintics. Macrocyclic lactone compounds have revolutionised the control of parasites in animals and are generally regarded as the most effective, least toxic and non-carcinogenic parasiticides yet developed. The M^{EC}WORMA & BOT paste is easy to administer under a wide range of conditions, with a smooth, sticky consistency that is well accepted. Abamectin has a wide safety margin at the recommended dose level especially in foals, pregnant mares, and aged or debilitated horses. There have been no adverse effects reported when used at double recommended doses in heavily parasitised horses.

DOSAGE AND ADMINISTRATION:

The recommended dose level of 200 µg/kg bodyweight M^{EC}WORMA & BOT controls the following parasites:

Large Strongyles: *Strongylus vulgaris* (adult & arterial larval stages), *Strongylus edentatus* (adult & tissue stages), *Strongylus equinus* (adult) and *Triodontophorus* spp. (adult). **Small Strongyles:** Including benzimidazole resistant strains (adult and immature) *Cyathostomum* spp., *Cylicoctylus* spp., *Cylicostephanus* spp., *Cylicodontophorus* spp. and *Gyalocephalus* spp. **Ascarids:** *Parascaris equorum* (adult & immature). **Pinworms:** *Oxyuris equi* (adult & immature). **Hairworms:** *Trichostrongylus axei* (adult). **Lungworms:** *Dictyocaulus arnfieldi* (adult and immature). **Bots:** *Gasterophilus* spp. (oral and stomach stages). **Intestinal Threadworms:** *Strongyloides westeri* (adult). **Neck Threadworms:** *Onchocerca* spp. (microfilariae) (cutaneous onchocerciasis). **Stomach Worms:** *Habronema muscae* (adult and cutaneous larval stage), *Draschia* spp. (cutaneous larval stage) - both cause **summer sores**.

A complete syringe treats a 600 kg body weight horse. The plunger is calibrated in 100 kg weight increments which each deliver 5.5 g of paste. Adjust the locking ring to the required body weight and depress plunger. Ensure paste is administered on the rear of the tongue and hold horse's head raised for a few seconds after dosing to ensure the paste is swallowed.

PARASITE CONTROL PROGRAM: A regular parasite control programme should be undertaken with particular attention paid to breeding stock, foals and yearlings. Contact International Animal Health Products or a veterinarian for a parasite control programme to suit your specific needs.

CAUTION (PROTECTION OF WILDLIFE): Free abamectin in M^{EC}WORMA & BOT may adversely affect fish and certain aquatic organisms. Do not contaminate dams, waterways or drains with this product or used container.

MEAT WITHHOLDING PERIOD: DO NOT USE less than 28 days before slaughter for human consumption.

POISONS SCHEDULE: S5.

REGULATORY STATUS: AUSTRALIAN PESTICIDES & VETERINARY MEDICINES AUTHORITY.

PACK SIZE: 33 g syringe (20 per carton); (50 per bucket).